

## 腫瘍関連、その他

1. Takahashi K, Togo Y, Saito K, Kiso H, Huang B, Tsukamoto H, Fujimura K, Bessho K. Two non-syndromic cases of multiple supernumerary teeth with different characteristics and a review of the literature. J Oral and Maxillofac Surg, Med, and pathol. 28: 250-254, 2016.
2. Idrees SR, Fujimura K, Bessho K. Dental trauma related to general anesthesia: should the anesthesiologist perform a preanesthetic dental evaluation? Oral Health Dent Manag. 13(2):271-4, 2014.
3. Fujimura K, Nishida M, Son A, Kunishima F, Segami N, Iizuka T, Kitaichi M. : Heterotopic calcification in advanced cervical lymph nodes with metastasis from squamous cell carcinoma of tongue: Report of two cases. Oral Oncology Extra. 40:117-122, 2004.
4. Bessho K, Fujimura K, Iizuka T. Experimental long-term study of titanium ions eluted from pure titanium miniplates. J Biomed Mater Res. 1995; 29(7):901-4.

## 顎変形症関連 （主に新しい外科矯正手術の開発について）

### IVSRO 下顎枝垂直矢状骨切り術

1. Fujimura K, Kazuhisa Bessho. : Rigid fixation of intraoral vertico-sagittal ramus osteotomy for mandibular prognathism. A Textbook of Advanced Oral and Maxillofacial Surgery. Chapter 27; Edited by Mohammad Hosein, Kalantar Motamedi. InTech. Croatia. pp 719-723. 2013.
2. Fujimura K, Bessho K. Rigid fixation of intraoral vertico-sagittal ramus osteotomy for mandibular prognathism. J Oral Maxillofac Surg. 2012; 70(5):1170-3.
3. Fujimura K, Segami N, Kobayashi S. Anatomical study of the complications of intraoral vertico-sagittal ramus osteotomy. J Oral Maxillofac Surg. 2006; 64(3):384-9.
4. Fujimura K, Segami N, Sato J, Kaneyama K, Nishimura M. : Comparison of the clinical outcomes of patients having sounds in the temporomandibular joint with skeletal mandibular deformities treated by vertico-sagittal ramus osteotomy or vertical ramus osteotomy. Oral Surg Oral Med Oral Pathol Oral Radiol Endod. 99(1):24-9, 2005.
5. Fujimura K, Segami N, Sato J, Kanayama K, Nishimura M, Demura N. Advantages of intraoral verticosagittal ramus osteotomy in skeletofacial deformity patients with temporomandibular joint disorders. J Oral Maxillofac Surg. 2004; 62(10):1246-52.
6. Nishimura M, Segami N, Sato J, Honjou M, Fujimura K. Transitional joint effusion in the mandibular prognathic surgery patient: intraoral vertical ramus osteotomy versus sagittal split ramus osteotomy. J Oral Maxillofac Surg. 2004; 62(5):545-8.

## 顎関節関連

1. Huang B, Takahashi K, Sakata T, Kiso H, Sugai M, Fujimura K, Shimizu A, Kosugi S, Sato T, Bessho K. Increased risk of temporomandibular joint closed lock: a case-control study of ANKH polymorphisms. *PLoS One*. 2011;6(10):e25503. doi:10.1371/journal.pone.0025503.
2. Koyama N, Okubo Y, Nakao K, Osawa K, Fujimura K, Bessho K. Pluripotency of mesenchymal cells derived from synovial fluid in patients with temporomandibular joint disorder. *Life Sci*. 2011; 89(19-20):741-7. doi:10.1016/j.lfs.2011.09.005.
3. Takafuji H, Suzuki T, Okubo Y, Fujimura K, Bessho K. Regeneration of articular cartilage defects in the temporomandibular joint of rabbits by fibroblast growth factor-2: a pilot study. *Int J Oral Maxillofac Surg*. 2007; 36(10):934-7.
4. Kaneyama K, Segami N, Sato J, Fujimura K, Nagao T, Yoshimura H. Prognostic factors in arthrocentesis of the temporomandibular joint: Comparison of bradykinin, leukotriene B4, prostaglandin E2, and substance P level in synovial fluid between successful and unsuccessful cases. *J Oral Maxillofac Surg*. 2007; 65(2):242-7.
5. Sato J, Segami N, Yoshitake Y, Kaneyama K, Yoshimura H, Fujimura K, Kitagawa Y. Specific expression of substance P in synovial tissues of patients with symptomatic, non-reducing internal derangement of the temporomandibular joint: comparison with clinical findings. *Br J Oral Maxillofac Surg*. 2007; 45(5):372-7.
6. Kaneyama K, Segami N, Shin-Ichi T, Fujimura K, Sato J, Nagao T. Anchored disc phenomenon with a normally positioned disc in the temporomandibular joint: characteristics and behaviour. *Br J Oral Maxillofac Surg*. 2007; 45(4):279-83.
7. Fujimura K, Segami N, Yoshitake Y, Tsuruoka N, Kaneyama K, Sato J, Kobayashi S. Electrophoretic separation of the synovial fluid proteins in patients with temporomandibular joint disorders. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod*. 2006; 101(4):463-8.
8. Sato J, Segami N, Yoshitake Y, Kaneyama K, Abe A, Yoshimura H, Fujimura K. Expression of capsaicin receptor TRPV-1 in synovial tissues of patients with symptomatic internal derangement of the temporomandibular joint and joint pain. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod*. 2005; 100(6):674-81.
9. Fujimura K, Kobayashi S, Yoshitake Y, Tsuruoka N, Kaneyama K, Segami N. Electrophoretically separation of the synovial fluid proteins in rabbit temporomandibular arthritis induced by mechanical loading. *J Oral Pathol Med*. 2005; 34(9):546-51.
10. Kaneyama K, Segami N, Sun W, Sato J, Fujimura K. Levels of soluble cytokine

factors in temporomandibular joint effusions seen on magnetic resonance images. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.* 2005; 99(4):411-8.

11. Kaneyama K, Segami N, Sun W, Sato J, Fujimura K. Analysis of tumor necrosis factor- $\alpha$ , interleukin-6, interleukin-1 $\beta$ , soluble tumor necrosis factor receptors I and II, interleukin-6 soluble receptor, interleukin-1 soluble receptor type II, interleukin-1 receptor antagonist, and protein in the synovial fluid of patients with temporomandibular joint disorders. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.* 2005; 99(3):276-84.

12. Sato J, Segami N, Kaneyama K, Mashiyama Y, Fujimura K, Yoshitake Y. Vascular endothelial growth factor concentrations in synovial fluids of patients with symptomatic internal derangement of the temporomandibular joint. *J Oral Pathol Med.* 2005; 34(3):170-7.

13. Fujimura K, Kobayashi S, Suzuki T, Segami N. Histologic evaluation of temporomandibular arthritis induced by mild mechanical loading in rabbits. *J Oral Pathol Med.* 2005; 34(3):157-63.

14. Fujimura K, Segami N, Sato J, Kaneyama K, Nishimura M. Comparison of the clinical outcomes of patients having sounds in the temporomandibular joint with skeletal mandibular deformities treated by vertico-sagittal ramus osteotomy or vertical ramus osteotomy. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.* 2005; 99(1):24-9.

15. Sato J, Segami N, Kaneyama K, Yoshimura H, Fujimura K, Yoshitake Y. Relationship of calcitonin gene-related peptide in synovial tissues and temporomandibular joint pain in humans. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.* 2004; 98(5):533-40.

16. Suzuki T, Bessho K, Fujimura K, Okubo Y, Segami N, Iizuka T. Regeneration of defects in the articular cartilage in rabbit temporomandibular joints by bone morphogenetic protein-2. *Br J Oral Maxillofac Surg.* 2002; 40(3):201-6.

17. Kaneyama K, Segami N, Nishimura M, Sato J, Suzuki T, Fujimura K. Osteoclastogenesis inhibitory factor/osteoprotegerin in synovial fluid from patients with temporomandibular disorders. *Int J Oral Maxillofac Surg.* 2003; 32(4):404-7.

18. Sato J, Segami N, Nishimura M, Suzuki T, Kaneyama K, Fujimura K. Clinical evaluation of arthroscopic eminoplasty for habitual dislocation of the temporomandibular joint: comparative study with conventional open eminectomy. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.* 2003; 95(4):390-5.

19. Kaneyama K, Segami N, Nishimura M, Sato J, Suzuki T, Fujimura K. Osteoclastogenesis inhibitory factor/osteoprotegerin in synovial fluid from patients with

temporomandibular disorders. *Int J Oral Maxillofac Surg.* 2003; 32(4):404-7.

20. Fujimura K, Bessho K, Okubo Y, Kusumoto K, Segami N, Iizuka T. The effect of fibroblast growth factor-2 on the osteoinductive activity of recombinant human bone morphogenetic protein-2 in rat muscle. *Arch Oral Biol.* 2002; 47(8):577-84.

21. Suzuki T, Bessho K, Fujimura K, Okubo Y, Segami N, Iizuka T. Regeneration of defects in the articular cartilage in rabbit temporomandibular joints by bone morphogenetic protein-2. *Br J Oral Maxillofac Surg.* 2002; 40(3):201-6.

22. Murakami K, Segami N, Fujimura K, Iizuka T. Correlation between pain and synovitis in patients with internal derangement of the temporomandibular joint. *J Oral Maxillofac Surg.* 1991; 49(11):1159-61.

23. Kaneyama K, Segami N, Nishimura M, Sato J, Fujimura K, Yoshimura H. The ideal lavage volume for removing bradykinin, interleukin-6, and protein from the temporomandibular joint by arthrocentesis. *J Oral Maxillofac Surg.* 2004; 62(6):657-61.

24. Nishimura M, Segami N, Kaneyama K, Sato J, Fujimura K. Comparison of cytokine level in synovial fluid between successful and unsuccessful cases in arthrocentesis of the temporomandibular joint. *J Oral Maxillofac Surg.* 2004; 62(3):284-7.

## 生体材料・再生医療関連

1. Isobe Y, Koyama N, Nakao K, Osawa K, Ikeno M, Yamanaka S, Okubo Y, Fujimura K, Bessho K. Comparison of human mesenchymal stem cells derived from bone marrow, synovial fluid, adult dental pulp, and exfoliated deciduous tooth pulp. *Int J Oral Maxillofac Surg.* 2015 Jul 30. pii: S0901-5027(15)00256-8. doi: 10.1016/j.ijom.2015.06.022.

2. Nakao K, Osawa K, Yasoda A, Yamanaka S, Fujii T, Kondo E, Koyama N, Kanamoto N, Miura M, Kuwahara K, Akiyama H, Bessho K, Nakao K.  
The Local CNP/GC-B system in growth plate is responsible for physiological endochondral bone growth. *Sci Rep* 5: 10554. 2015.

3. Koyama N, Kondo S, Watanabe T, Nishioka M, Kaizaki Y, Kawai Y.  
A case of IgG4-related disease founded from sclerosing sialadenitis.  
*J Oral Maxillofac Surg Med Path* 27(6): 812-6. 2015

4. Koyama N, Miura M, Nakao K, Kondo E, Fujii T, Taura D, Kanamoto N, Sone M, Yasoda A, Arai H, Bessho K, Nakao K.  
Human induced pluripotent stem cells differentiated into chondrogenic lineage via generation of mesenchymal progenitor cells.  
*Stem Cells Dev* 22(1): 102-13. 2013.

5. Nakao K, Okubo Y, Yasoda A, Koyama N, Osawa K, Isobe Y, Kondo E, Fujii T, Miura M, Nakao K, Bessho K.  
The effects of C-type natriuretic peptide on craniofacial skeletogenesis.  
J Dent Res 92(1): 58-64. 2013
6. Osawa K, Okubo Y, Nakao K, Koyama N, Bessho K.  
Feasibility of BMP-2 Gene Therapy Using an Ultra-Fine Needle.  
Chapter 9. Targets in Gene Therapy, Yongping You (Ed.), ISBN: 978-953-307-540-2, InTech, p159-166, 2011
7. Koyama N, Okubo Y, Nakao K, Bessho K.  
Evaluation of Pluripotency in Human Dental Pulp Cells.  
J Oral Maxillofac Surg 67(3): 501-506. 2009
8. Osawa K, Okubo Y, Nakao K, Koyama N, Bessho K.  
Osteoinduction by microbubble-enhanced transcutaneous sonoporation of human bone morphogenetic protein-2.  
J Gene Med 11(7): 633-641. 2009
9. Osawa K, Okubo Y, Nakao K, Koyama N, Bessho K.  
Osteoinduction by repeat plasmid injection of human bone morphogenetic protein-2.  
J Gene Med 12(12): 937-44. 2010
10. Koyama N, Okubo Y, Nakao K, Osawa K, Bessho K.  
Experimental study of osteoinduction using a new material as a carrier for bone morphogenetic protein-2.  
Br J Oral Maxillofac Surg 49(4): 314-8. 2011
11. Okubo Y, Reddi AH, Nakao K, Koyama N, Osawa K, Bessho K.  
Terminal Differentiation of Chondrocytes is Stimulated by Thyroxine.  
Current Topics in Biochemical Research, Research Signpost, Trivandrum, p57-62, 2007.
12. Kusumoto K, Bessho K, Fujimura K, Okubo Y, Wang Y, Kakudo N, Ogawa Y. Tissue reaction at the implantation of recombinant human bone morphogenetic protein-2 into the skeletal muscle. Int J Tissue React. 2005; 27(4):181-8.
13. Fujimura K, Bessho K, Okubo Y, Segami N, Iizuka T. A bioactive bone cement containing Bis-GMA resin and A-W glass-ceramic as an augmentation graft material on mandibular bone. Clin Oral Implants Res. 2003; 14(5):659-67.
14. Okubo Y, Bessho K, Fujimura K, Kusumoto K, Suzuki T, Segami N, Ogawa Y, Iizuka T. Preclinical study of recombinant human bone morphogenetic protein-2: application of hyperbaric oxygenation during bone formation under unfavourable condition. Int J Oral Maxillofac Surg. 2003; 32(3):313-7.
15. Fujimura K, Bessho K, Okubo Y, Kusumoto K, Segami N, Iizuka T. The effect of fibroblast growth factor-2 on the osteoinductive activity of recombinant human

bone morphogenetic protein-2 in rat muscle. *Arch Oral Biol.* 2002; 47(8):577-84.

16. Kusumoto K, Bessho K, Fujimura K, Akioka J, Okubo Y, Wang Y, Iizuka T, Ogawa Y. Osteoinduction by recombinant human bone morphogenetic protein-2 in muscles of non-human primates. *J Int Med Res.* 2002; 30(3):251-9.

17. Okubo Y, Bessho K, Fujimura K, Kaihara S, Iizuka T, Miyatake S. The time course study of osteoinduction by bone morphogenetic protein-2 via adenoviral vector. *Life Sci.* 2001; 70(3):325-36.

18. Okubo Y, Bessho K, Fujimura K, Kusumoto K, Ogawa Y, Iizuka T. Expression of bone morphogenetic protein in the course of osteoinduction by recombinant human bone morphogenetic protein-2. *Clin Oral Implants Res.* 2002; 13(1):80-5.

19. Fujimura K, Bessho K, Kusumoto K, Konishi Y, Ogawa Y, Iizuka T. Experimental osteoinduction by recombinant human bone morphogenetic protein 2 in tissue with low blood flow: a study in rats. *Br J Oral Maxillofac Surg.* 2001; 39(4):294-300.

20. Okubo Y, Bessho K, Fujimura K, Iizuka T, Miyatake SI. In vitro and in vivo studies of a bone morphogenetic protein-2 expressing adenoviral vector. *J Bone Joint Surg Am.* 2001; 83-A Suppl 1(Pt 2):S99-104.

21. Okubo Y, Bessho K, Fujimura K, Kusumoto K, Ogawa Y, Iizuka T. Effect of hyperbaric oxygenation on bone induced by recombinant human bone morphogenetic protein-2. *Br J Oral Maxillofac Surg.* 2001; 39(2):91-5.

22. Bessho K, Konishi Y, Kaihara S, Fujimura K, Okubo Y, Iizuka T. Bone induction by *Escherichia coli*-derived recombinant human bone morphogenetic protein-2 compared with Chinese hamster ovary cell-derived recombinant human bone morphogenetic protein-2. *Br J Oral Maxillofac Surg.* 2000; 38(6):645-649.

23. Kusumoto K, Bessho K, Fujimura K, Akioka J, Okubo Y, Ogawa Y, Iizuka T. The effect of blood supply in muscle and an elevated muscle flap on endogenous tissue-engineered bone by rhBMP-2 in the rat. *Ann Plast Surg.* 2000; 45(4):408-14.

24. Okubo Y, Bessho K, Fujimura K, Kusumoto K, Ogawa Y, Iizuka T. Osteogenesis by recombinant human bone morphogenetic protein-2 at skeletal sites. *Clin Orthop Relat Res.* 2000; (375):295-301.

25. Okubo Y, Bessho K, Fujimura K, Kusumoto K, Ogawa Y, Iizuka T. Effect of elcatonin on osteoinduction by recombinant human bone morphogenetic protein-2. *Biochem Biophys Res Commun.* 2000; 269(2):317-21.

26. Okubo Y, Bessho K, Fujimura K, Konishi Y, Kusumoto K, Ogawa Y, Iizuka T. Osteoinduction by recombinant human bone morphogenetic protein-2 at intramuscular, intermuscular, subcutaneous and intrafatty sites. *Int J Oral Maxillofac Surg.* 2000; 29(1):62-6.

27. Okubo Y, Bessho K, Fujimura K, Iizuka T, Miyatake SI. Osteoinduction by bone morphogenetic protein-2 via adenoviral vector under transient immunosuppression. *Biochem Biophys Res Commun.* 2000; 267(1):382-7.
28. Yoshida K, Bessho K, Fujimura K, Konishi Y, Kusumoto K, Ogawa Y, Iizuka T. Enhancement by recombinant human bone morphogenetic protein-2 of bone formation by means of porous hydroxyapatite in mandibular bone defects. *J Dent Res.* 1999; 78(9):1505-10.
29. Okubo Y, Bessho K, Fujimura K, Iizuka T, Miyatake S. Expression of bone morphogenetic protein-2 via adenoviral vector in C2C12 myoblasts induces differentiation into the osteoblast lineage. *Biochem Biophys Res Commun.* 1999; 262(3):739-43.
30. Bessho K, Kusumoto K, Fujimura K, Konishi Y, Ogawa Y, Tani Y, Iizuka T. Comparison of recombinant and purified human bone morphogenetic protein. *Br J Oral Maxillofac Surg.* 1999; 37(1):2-5.
31. Okubo Y, Bessho K, Fujimura K, Kusumoto K, Ogawa Y, Tani Y, Iizuka T. Comparative study of intramuscular and intraskeletal osteogenesis by recombinant human bone morphogenetic protein-2. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.* 1999; 87(1):34-8.
32. Kusumoto K, Bessho K, Fujimura K, Akioka J, Ogawa Y, Iizuka T. Prefabricated muscle flap including bone induced by recombinant human bone morphogenetic protein-2: an experimental study of ectopic osteoinduction in a rat latissimus dorsi muscle flap. *Br J Plast Surg.* 1998; 51(4):275-80.
33. Yoshida K, Bessho K, Fujimura K, Kusumoto K, Ogawa Y, Tani Y, Iizuka T. Osteoinduction capability of recombinant human bone morphogenetic protein-2 in intramuscular and subcutaneous sites: an experimental study. *J Craniomaxillofac Surg.* 1998; 26(2):112-5.
34. Kusumoto K, Bessho K, Fujimura K, Ogawa Y, Iizuka T. Intramuscular osteoinduction and bone marrow formation by the implantation of rhBMP-2 with atelopeptide type I collagen. *Br J Oral Maxillofac Surg.* 1997 Dec;35(6):433-7.
35. Kusumoto K, Bessho K, Fujimura K, Akioka J, Ogawa Y, Iizuka T. Comparison of ectopic osteoinduction in vivo by recombinant human BMP-2 and recombinant Xenopus BMP-4/7 heterodimer. *Biochem Biophys Res Commun.* 1997; 239(2):575-9.
36. Kusumoto K, Bessho K, Fujimura K, Konishi Y, Ogawa Y, Iizuka T. Self-regenerating bone implant: ectopic osteoinduction following intramuscular implantation of a combination of rhBMP-2, atelopeptide type I collagen and porous hydroxyapatite. *J Craniomaxillofac Surg.* 1996; 24(6):360-5.

37. Kusumoto K, Bessho K, Fujimura K, Konishi Y, Ogawa Y, Iizuka T. Comparative study of bone marrow induced by purified BMP and recombinant human BMP-2. *Biochem Biophys Res Commun.* 1995; 215(1):205-11.
38. Fujimura K, Bessho K, Kusumoto K, Ogawa Y, Iizuka T. Experimental studies on bone inducing activity of composites of atelopeptide type I collagen as a carrier for ectopic osteoinduction by rhBMP-2. *Biochem Biophys Res Commun.* 1995; 208(1):316-22.